

WHAT IS CLAIMED IS:

1. A method for treating disorders associated with radical depolarization of excitable membranes by activating a K<sub>ATP</sub> channel which comprises administering to an individual in need thereof an effective amount of an active agent selected from the group consisting of:

- 5 (a) a compound of the following formula

Cys-Xaa<sub>1</sub>-Ile-Xaa<sub>2</sub>-Asn-Gln-Xaa<sub>3</sub>-Cys-Xaa<sub>4</sub>-Gln-Xaa<sub>5</sub>-Leu-Asp-Asp-Cys-Cys-Ser-Xaa<sub>1</sub>-Xaa<sub>3</sub>-Cys-Asn-Xaa<sub>1</sub>-Xaa<sub>4</sub>-Asn-Xaa<sub>3</sub>-Cys-Val (SEQ ID NO:1), wherein Xaa<sub>1</sub> and Xaa<sub>3</sub> are independently Arg, homoarginine, ornithine, Lys, N-methyl-Lys, N,N-dimethyl-Lys, N,N,N-trimethyl-Lys, any synthetic basic amino acid, His or halo-His; Xaa<sub>2</sub> is Pro or hydroxy-Pro (Hyp); Xaa<sub>4</sub> is Phe, Tyr, meta-Tyr, ortho-Tyr, nor-Tyr, mono-halo-Tyr, di-halo-Tyr, O-sulpho-Tyr, O-phospho-Tyr, nitro-Tyr, Trp (D or L), neo-Trp, halo-Trp (D or L) or any synthetic aromatic amino acid; and Xaa<sub>5</sub> is His or halo-His,

- (b) an analog of the compound of (a), said analog selected from the group consisting of:

15 κ-PVIIA[R18A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Ala-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:2);

κ-PVIIA[R22A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Ala-Phe-Asn-Lys-Cys-Val (SEQ ID NO:3);

20 κ-PVIIA[I3A]: Cys-Arg-Ala-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:4);

κ-PVIIA[K19A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Ala-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:5);

κ-PVIIA[R2A]: Cys-Ala-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:6);

25 κ-PVIIA[F9A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Ala-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:7);

κ-PVIIA[K25A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Ala-Cys-Val (SEQ ID NO:8);

κ-PVIIA[R2K]: Cys-Lys-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:9);

5

10  
15  
20  
25  
30

20

25

30

- $\kappa$ -PVIIA[K7A]: Cys-Arg-Ile-Hyp-Asn-Gln-Ala-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:10);
- $\kappa$ -PVIIA[F9M]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Met-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:11);
- 5  $\kappa$ -PVIIA[F9Y]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Tyr-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:12);
- $\kappa$ -PVIIA[R2Q]: Cys-Gln-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:13);
- 10  $\kappa$ -PVIIA[H11A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-Ala-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:14);
- $\kappa$ -PVIIA[D14A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Ala-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:15);
- $\kappa$ -PVIIA[Q6A]: Cys-Arg-Ile-Hyp-Asn-Ala-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:16);
- $\kappa$ -PVIIA[N21A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Ala-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:17);
- $\kappa$ -PVIIA[S17A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ala-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:18);
- 20  $\kappa$ -PVIIA[N24A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Ala-Lys-Cys-Val (SEQ ID NO:19);
- $\kappa$ -PVIIA[L12A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Ala-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:20);
- $\kappa$ -PVIIA[D13A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Ala-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:21);
- 25  $\kappa$ -PVIIA[Q10A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Ala-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:22);
- $\kappa$ -PVIIA[V27A]: Cys-Arg-Ile-Hyp-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Ala (SEQ ID NO:23);
- 30  $\kappa$ -PVIIA[O4A]: Cys-Arg-Ile-Ala-Asn-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:24); and
- $\kappa$ -PVIIA[N5A]: Cys-Arg-Ile-Hyp-Ala-Gln-Lys-Cys-Phe-Gln-His-Leu-Asp-Asp-Cys-Cys-Ser-Arg-Lys-Cys-Asn-Arg-Phe-Asn-Lys-Cys-Val (SEQ ID NO:25);

- (c) a derivative of (a) or (b); and
- (d) a physiologically acceptable salt thereof.

2. The method of claim 1, wherein Xaa<sub>2</sub> is hydroxy-Pro.
3. The method of claim 1, wherein Xaa<sub>1</sub> is Arg, Xaa<sub>3</sub> is Lys, Xaa<sub>4</sub> is Phe and Xaa<sub>5</sub> is His.
4. The method of claim 3, wherein Xaa<sub>2</sub> is hydroxy-Pro.
5. The method of claim 1, wherein said disorder is cardiac ischemia.
6. The method of claim 1, wherein said disorder is cerebral ischemia.
7. The method of claim 1, wherein said disorder is asthma.
8. The method of claim 1, wherein said disorder is ocular ischemia.
9. The method of claim 1, wherein the derivative is peptide of (a) or (b) in which the Arg residues may be substituted by Lys, ornithine, homoarginine, nor-Lys, N-methyl-Lys, N,N-dimethyl-Lys, N,N,N-trimethyl-Lys or any synthetic basic amino acid; the Lys residues may be substituted by Arg, ornithine, homoarginine, nor-Lys, or any synthetic basic amino acid; the Tyr residues may be substituted with any synthetic hydroxy containing amino acid; the Ser residues may be substituted with Thr or any synthetic hydroxylated amino acid; the Thr residues may be substituted with Ser or any synthetic hydroxylated amino acid; the Phe and Trp residues may be substituted with any synthetic aromatic amino acid; the Asn, Ser, Thr or Hyp residues may be glycosylated (contain an N-glycan or an O-glycan); the Cys residues may be in D or L configuration and may optionally be substituted with homocysteine (D or L); the Tyr residues may also be substituted with the 3-hydroxyl or 2-hydroxyl isomers (meta-Tyr or ortho-Tyr, respectively) and corresponding O-sulpho- and O-phospho-derivatives; the acidic amino acid residues may be substituted with any synthetic acidic amino acid, e.g., tetrazolyl derivatives of Gly and Ala; the aliphatic amino acids may be substituted by synthetic derivatives bearing non-natural aliphatic branched or linear side

chains  $C_nH_{2n+2}$  up to and including  $n=8$ ; and pairs of Cys residues may be replaced pairwise with isoteric lactam or ester-thioether replacements, such as Ser/(Glu or Asp), Lys/(Glu or Asp) or Cys/Ala combinations.

ADD  
β1